

COMMUNICABLE DISEASE CENTER

Morbidity and Mortality



Vol. 15, No. 2

WEEKLY
REPORTWeek Ending
January 15, 1966

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE

PUBLIC HEALTH SERVICE

INFECTIOUS HEPATITIS

NASSAU COUNTY, LONG ISLAND, NEW YORK

Between September and December 1965, there were 90 confirmed cases of infectious hepatitis in the Elmont-North Valley Stream area of Nassau County, Long Island, New York. The distribution of cases in time, by date of onset, is shown in Figure 1. There is a sharply rising and falling curve with most of the cases occurring between the middle of October and the middle of November over a period of approximately 5 weeks. In Table 1 the age and sex distributions of cases are listed. Although a predominance of the cases was among teen-agers, there was an unusual occurrence among adults. Twenty of the 90 cases were 40 years of age and older.

Infectious Hepatitis - Nassau County, Long Island, New York	9
Reported Cases of Infectious Syphilis	11
Hepatitis Surveillance Summary	16

CONTENTS

The clustering of cases over a short period of time and the adult age distribution of cases both strongly suggested a common-source exposure. The water supply to the area where the cases lived was common to other more extensive areas in which cases of infectious hepatitis were not known to have occurred during the epidemic period. Twenty different dairies were named by patients

(Continued on page 10)

CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES
(Cumulative totals include revised and delayed reports through previous weeks)

DISEASE	2nd WEEK ENDED		MEDIAN 1961-1965	CUMULATIVE, FIRST 2 WEEKS		
	JANUARY 15, 1966	JANUARY 16, 1965		1966	1965	MEDIAN 1961-1965
Aseptic meningitis	41	30	30	71	72	47
Brucellosis	4	2	4	4	10	10
Diphtheria	2	6	9	3	9	12
Encephalitis, primary:						
Arthropod-borne & unspecified	24	37	---	43	70	---
Encephalitis, post-infectious	15	19	---	24	24	---
Hepatitis, serum	22	795	1,093	1,361	1,483	1,784
Hepatitis, infectious	695					
Measles (rubeola)	5,544	6,746	7,589	9,405	11,949	12,497
Poliomyelitis, Total (including unspecified)	—	—	6	1	—	7
Paralytic	—	—	4	—	—	6
Nonparalytic	—	—	---	---	---	---
Meningococcal infections, Total	94	74	56	157	126	98
Civilian	87	72	---	145	124	---
Military	7	2	---	12	2	---
Rubella (German measles)	930	---	---	1,326	---	---
Streptococcal sore throat & Scarlet fever	9,405	9,974	8,457	16,766	18,654	15,544
Tetanus	—	5	---	3	8	---
Tularemia	4	9	---	8	15	---
Typhoid fever	4	3	3	7	5	7
Typhus, tick-borne (Rky. Mt. Spotted fever)	2	2	---	7	2	---
Rabies in Animals	67	88	64	139	180	114

NOTIFIABLE DISEASES OF LOW FREQUENCY

	Cum.		Cum.
Anthrax	—	Botulism	—
Leptospirosis	—	Trichinosis: N.Y. Up-State-5, N.J.-1	10
Malaria: Pa.-1, Va.-1, Oreg.-1	8	Rabies in Man	—
Psittacosis: N.C.-1	3	Rubella, Congenital Syndrome: Pa.-1	1
Typhus, murine	—		

INFECTIOUS HEPATITIS **NASSAU COUNTY, LONG ISLAND, NEW YORK**

(Continued from front page)

as sources of milk supply, and raw shellfish were consumed by only 15 of the 90 patients in the 2 months preceding their respective illnesses. When the patients were questioned as to where they purchased cold cuts or freshly prepared salads within this time, they named nine various delicatessens and groceries. Of the 76 cases with dates of onset between October 13 and November 21, 66 (86.8 percent) had purchased these foods from one local delicatessen in the preceding 2 months. A control group of 94 households was randomly selected from the same neighborhoods as the cases. Fifteen of the 94 control families, or 16.0 percent, had patronized the suspect delicatessen within the preceding 2 months.

The delicatessen involved was a small, privately-owned store which was run by the owner and two employees. The owner of the store personally prepared the various salads, sliced the cold cuts and served customers from behind the counter. He had had an illness indistinguishable from infectious hepatitis on both clinical and laboratory grounds with onset sometime during the last week of September. However, he continued to work in the store through October 16 despite the occurrence of jaundice, documented on October 13, as shown in Figure 1. It seemed likely that contamination of food had occurred at this delicatessen, but no single food item was incriminated.

None of the five cases subsequent to November 21 had a history of purchases from this delicatessen, but all

Table 1
AGE AND SEX DISTRIBUTION FOR 90 CASES
OF INFECTIOUS HEPATITIS
 Elmont-North Valley Stream Area, Nassau County, New York
 September-December 1965


Age	Male	Female	Total
0-4	—	—	—
5-9	6	2	8
10-14	8	12	20
15-19	11	9	20
20-24	7	—	7
25-29	2	1	3
30-34	3	2	5
35-39	5	2	7
40-44	4	8	12
45+	7	1	8
Total	53	37	90


of the five gave a history of personal contact with a previous known case.

(Reported by Dr. I. Jackson Tartakow, Deputy Commissioner, Nassau County Health Department, Mineola Long Island, New York; Dr. Julia Freitag, Acting Director, Office of Epidemiology, New York State Department of Health, Albany, New York; and an EIS Officer.)

Figure 1.

DATES OF ONSET OF 90 CASES OF INFECTIOUS HEPATITIS **ELMONT-NORTH VALLEY STREAM AREA, NASSAU COUNTY, NEW YORK** **SEPTEMBER-DECEMBER 1965**





DECEMBER 1965 AND DECEMBER 1964

Reporting Area	December		Cumulative Jan. - Dec.		Reporting Area	December		Cumulative Jan. - Dec.	
	1965	1964	1965	1964		1965	1964	1965	1964
NEW ENGLAND.....	41	42	480	483	EAST SOUTH CENTRAL.....	196	171	2,676	1,922
Maine.....	-	2	2	7	Kentucky.....	12	20	138	178
New Hampshire.....	1	1	28	10	Tennessee.....	33	38	504	479
Vermont.....	-	-	4	3	Alabama.....	39	78	1,421	902
Massachusetts.....	33	21	302	284	Mississippi.....	52	35	613	36
Rhode Island.....	-	1	21	19	WEST SOUTH CENTRAL.....	188	182	2,331	2,599
Connecticut.....	7	17	123	160	Arkansas.....	4	22	202	227
MIDDLE ATLANTIC.....	467	470	4,901	5,428	Louisiana.....	54	55	660	734
Update New York.....	43	42	549	632	Oklahoma.....	16	111	151	150
New York City.....	28	299	2,853	3,165	Texas.....	120	89	1,358	1,488
Pa. (Excl. Phila.).....	18	16	189	182	MOUNTAIN.....	41	35	545	539
Philadelphia.....	59	22	365	309	Montana.....	3	1	22	32
New Jersey.....	52	91	945	1,140	Idaho.....	-	1	9	8
EAST NORTH CENTRAL.....	277	249	3,036	2,474	Wyoming.....	-	-	9	8
Ohio.....	64	58	648	584	Colorado.....	6	2	51	36
Indiana.....	14	63	70	63	New Mexico.....	6	10	112	180
Downstate Illinois.....	12	22	199	161	Arizona.....	21	14	266	223
Chicago.....	92	103	1,186	974	Utah.....	4	1	21	14
Michigan.....	92	58	862	622	Nevada.....	1	6	55	39
Wisconsin.....	3	3	78	63	PACIFIC.....	179	178	2,031	2,213
WEST NORTH CENTRAL.....	45	44	550	550	Washington.....	1	2	69	73
Minnesota.....	4	9	124	99	Oregon.....	11	5	49	73
Iowa.....	6	4	59	37	California.....	160	167	1,873	2,039
Missouri.....	25	19	248	252	Alaska.....	1	-	8	9
North Dakota.....	-	-	2	-	Hawaii.....	1	4	32	19
South Dakota.....	1	5	39	53	U. S. TOTAL.....	1,937	1,814	23,290	22,969
Nebraska.....	5	4	86	56	TERRITORIES.....	85	73	889	878
Kansas.....	4	3	36	28	Puerto Rico.....	85	71	865	849
SOUTH ATLANTIC.....	503	443	6,740	6,761	Virgin Islands.....	-	2	24	29
Delaware.....	5	3	56	84					
Maryland.....	42	38	489	514					
District of Columbia.....	47	36	493	693					
Virginia.....	33	20	327	299					
West Virginia.....	12	8	107	59					
North Carolina.....	66	77	1,004	1,109					
South Carolina.....	79	58	851	860					
Georgia.....	61	83	1,057	1,139					
Florida.....	158	120	2,356	2,004					

Note: Cumulative Totals include revised and delayed reports through previous months.

Note: Cumulative Totals include revised and delayed reports through previous months.

JANUARY 15, 1966 AND JANUARY 16, 1965 (2nd WEEK)

[illegible]

Morbidity and Mortality Weekly Report

13

CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES
FOR WEEKS ENDED
JANUARY 15, 1966 AND JANUARY 16, 1965 (2nd WEEK) - Continued

AREA	MEASLES (Rubeola)			MENINGOCOCCAL INFECTIONS, TOTAL			POLIO MYELITIS				RUBELLA
	1966	Cumulative		1966	Cumulative		Total		Paralytic		
		1966	1965		1966	1965	1966	1965	1966	Cumulative 1966	
UNITED STATES...	5,544	9,405	11,949	94	157	126	-	-	-	-	930
NEW ENGLAND.....	76	122	3,747	7	12	7	-	-	-	-	61
Maine.....	9	16	495	-	-	2	-	-	-	-	8
New Hampshire.....	2	3	87	3	3	-	-	-	-	-	-
Vermont.....	30	47	20	-	1	-	-	-	-	-	-
Massachusetts.....	22	28	1,966	3	4	4	-	-	-	-	20
Rhode Island.....	3	9	362	1	1	1	-	-	-	-	3
Connecticut.....	10	19	817	-	3	-	-	-	-	-	30
MIDDLE ATLANTIC.....	775	1,617	422	24	35	19	-	-	-	-	167
New York City.....	449	740	67	9	11	4	-	-	-	-	26
New York, Up-State.....	75	259	157	3	5	5	-	-	-	-	141
New Jersey.....	51	241	54	5	10	6	-	-	-	-	-
Pennsylvania.....	200	377	144	7	9	4	-	-	-	-	-
EAST NORTH CENTRAL...	2,835	4,101	1,731	14	24	16	-	-	-	-	313
Ohio.....	93	212	297	6	10	8	-	-	-	-	-
Indiana.....	172	221	53	1	3	-	-	-	-	-	57
Illinois.....	853	978	56	2	3	6	-	-	-	-	81
Michigan.....	292	488	943	4	6	-	-	-	-	-	53
Wisconsin.....	1,425	2,202	382	1	2	2	-	-	-	-	122
WEST NORTH CENTRAL...	92	231	825	3	8	8	-	-	-	-	50
Minnesota.....	1	66	8	1	3	-	-	-	-	-	7
Iowa.....	44	68	430	-	1	-	-	-	-	-	34
Missouri.....	5	16	82	1	1	4	-	-	-	-	2
North Dakota.....	41	76	277	-	-	3	-	-	-	-	7
South Dakota.....	-	1	16	1	1	-	-	-	-	-	-
Nebraska.....	1	4	12	-	-	-	-	-	-	-	-
Kansas.....	NN	NN	NN	-	2	1	-	-	-	-	-
SOUTH ATLANTIC.....	445	965	1,632	13	26	25	-	-	-	-	18
Delaware.....	10	17	51	-	-	1	-	-	-	-	-
Maryland.....	68	125	18	3	4	-	-	-	-	-	5
Dist. of Columbia..	15	32	-	-	-	-	-	-	-	-	-
Virginia.....	10	20	156	-	-	3	-	-	-	-	5
West Virginia.....	250	559	1,266	-	-	2	-	-	-	-	8
North Carolina.....	3	13	50	2	4	2	-	-	-	-	-
South Carolina.....	44	72	17	5	7	-	-	-	-	-	-
Georgia.....	10	20	11	-	4	9	-	-	-	-	-
Florida.....	35	107	63	3	7	8	-	-	-	-	-
EAST SOUTH CENTRAL...	571	1,007	559	1	1	11	-	-	-	-	58
Kentucky.....	214	282	10	-	-	7	-	-	-	-	17
Tennessee.....	336	687	411	1	1	2	-	-	-	-	40
Alabama.....	9	9	62	-	-	2	-	-	-	-	1
Mississippi.....	12	29	76	-	-	-	-	-	-	-	-
WEST SOUTH CENTRAL...	253	492	852	6	10	16	-	-	-	-	-
Arkansas.....	16	18	2	1	1	2	-	-	-	-	-
Louisiana.....	9	10	1	2	2	4	-	-	-	-	-
Oklahoma.....	1	1	5	-	-	3	-	-	-	-	-
Texas.....	227	463	844	3	7	7	-	-	-	-	-
MOUNTAIN.....	249	408	1,288	8	8	-	-	-	-	-	95
Montana.....	47	79	468	-	-	-	-	-	-	-	35
Idaho.....	99	107	196	-	-	-	-	-	-	-	3
Wyoming.....	1	7	54	-	-	-	-	-	-	-	-
Colorado.....	22	27	181	8	8	-	-	-	-	-	2
New Mexico.....	-	2	47	-	-	-	-	-	-	-	-
Arizona.....	77	175	30	-	-	-	-	-	-	-	53
Utah.....	3	7	309	-	-	-	-	-	-	-	2
Nevada.....	-	4	3	-	-	-	-	-	-	-	-
PACIFIC.....	248	462	893	18	33	24	-	-	-	-	168
Washington.....	56	132	141	-	1	-	-	-	-	-	88
Oregon.....	31	65	304	1	2	1	-	-	-	-	33
California.....	156	258	349	9	21	22	-	-	-	-	43
Alaska.....	-	-	9	7	7	1	-	-	-	-	-
Hawaii.....	5	7	90	1	2	-	-	-	-	-	4
Puerto Rico.....	59	108	58	-	-	-	-	-	-	-	2

JANUARY 15, 1966 AND JANUARY 16, 1965 (2nd WEEK) - Continued

[illegible]

Week No. Table 4. DEATHS IN 122 UNITED STATES CITIES FOR WEEK ENDED JANUARY 15, 1966

2 (By place of occurrence and week of filing certificate. Excludes fetal deaths)									
Area	All Causes		Pneumonia and Influenza All Ages	Under 1 year All Causes	Area	All Causes		Pneumonia and Influenza All Ages	Under 1 year All Causes
	All Ages	65 years and over				All Ages	65 years and over		
NEW ENGLAND:	863	525	43	34	SOUTH ATLANTIC:	1,262	654	68	91
Boston, Mass.-----	279	155	6	9	Atlanta, Ga.-----	172	71	5	16
Bridgeport, Conn.-----	47	33	4	1	Baltimore, Md.-----	292	160	9	23
Cambridge, Mass.-----	34	20	3	1	Charlotte, N. C.-----	40	19	2	2
Fall River, Mass.-----	38	28	1	3	Jacksonville, Fla.-----	75	34	9	6
Hartford, Conn.-----	71	31	3	7	Miami, Fla.-----	74	50	1	2
Lowell, Mass.-----	31	23	3	1	Norfolk, Va.-----	71	37	7	2
Lynn, Mass.-----	24	15	1	1	Richmond, Va.-----	78	34	1	13
New Bedford, Mass.-----	33	28	4	1	Savannah, Ga.-----	39	16	5	2
New Haven, Conn.-----	68	43	1	1	St. Petersburg, Fla.---	77	65	5	-
Providence, R. I.-----	64	32	3	4	Tampa, Fla.-----	88	58	10	4
Somerville, Mass.-----	23	18	4	-	Washington, D. C.-----	214	88	12	16
Springfield, Mass.-----	58	40	6	1	Wilmington, Del.-----	42	22	2	5
Waterbury, Conn.-----	24	11	-	1					
Worcester, Mass.-----	69	48	4	4					
MIDDLE ATLANTIC:	3,583	2,083	176	165	EAST SOUTH CENTRAL:	758	396	42	39
Albany, N.Y.-----	65	27	3	3	Birmingham, Ala.-----	131	65	3	7
Allentown, Pa.-----	33	23	2	-	Chattanooga, Tenn.-----	76	39	10	1
Buffalo, N. Y.-----	164	97	8	12	Knoxville, Tenn.-----	70	44	6	1
Camden, N. J.-----	46	26	2	2	Louisville, Ky.-----	143	81	15	6
Elizabeth, N. J.-----	41	20	1	3	Memphis, Tenn.-----	146	75	2	12
Erie, Pa.-----	43	30	1	3	Mobile, Ala.-----	45	22	-	7
Jersey City, N. J.-----	82	46	6	3	Montgomery, Ala.-----	43	16	4	4
Newark, N. J.-----	117	52	6	10	Nashville, Tenn.-----	104	54	2	1
New York City, N. Y.---	1,832	1,063	86	77	WEST SOUTH CENTRAL:	1,191	618	56	89
Paterson, N. J.-----	47	24	4	3	Austin, Tex.-----	46	25	5	1
Philadelphia, Pa.-----	515	313	14	21	Baton Rouge, La.-----	14	8	2	-
Pittsburgh, Pa.-----	195	113	4	10	Corpus Christi, Tex.---	35	14	-	3
Reading, Pa.-----	50	26	5	4	Dallas, Tex.-----	154	80	3	13
Rochester, N. Y.-----	109	71	10	2	El Paso, Tex.-----	37	14	4	2
Schenectady, N. Y.-----	30	22	-	-	Fort Worth, Tex.-----	96	61	4	6
Scranton, Pa.-----	29	18	3	-	Houston, Tex.-----	231	111	17	17
Syracuse, N. Y.-----	57	34	3	3	Little Rock, Ark.-----	72	41	3	5
Trenton, N. J.-----	69	38	9	4	New Orleans, La.-----	153	75	6	14
Utica, N. Y.-----	25	20	6	2	Oklahoma City, Okla.---	102	53	4	9
Yonkers, N. Y.-----	34	20	3	3	San Antonio, Tex.-----	128	73	4	6
					Shreveport, La.-----	50	20	5	9
					Tulsa, Okla.-----	73	43	5	4
EAST NORTH CENTRAL:	2,693	1,560	130	143	MOUNTAIN:	438	260	21	21
Akron, Ohio-----	66	46	-	3	Albuquerque, N. Mex.---	46	21	1	5
Canton, Ohio-----	30	24	3	-	Colorado Springs, Colo.	18	13	3	1
Chicago, Ill.-----	845	433	40	47	Denver, Colo.-----	133	81	11	4
Cincinnati, Ohio-----	190	113	2	9	Ogden, Utah-----	21	14	1	-
Cleveland, Ohio-----	248	154	7	14	Phoenix, Ariz.-----	106	65	3	5
Columbus, Ohio-----	106	72	4	3	Pueblo, Colo.-----	17	11	-	-
Dayton, Ohio-----	67	42	13	3	Salt Lake City, Utah--	47	26	1	5
Detroit, Mich.-----	352	207	18	25	Tucson, Ariz.-----	50	29	1	1
Evansville, Ind.-----	49	32	6	2					
Flint, Mich.-----	44	23	2	3	PACIFIC:	1,851	1,108	74	90
Fort Wayne, Ind.-----	37	24	6	-	Berkeley, Calif.*-----	20	13	-	1
Cary, Ind.-----	37	19	4	2	Fresno, Calif.-----	63	33	4	5
Grand Rapids, Mich.---	33	18	-	1	Glendale, Calif.*-----	45	32	1	1
Indianapolis, Ind.-----	147	83	8	1	Honolulu, Hawaii-----	51	27	4	5
Madison, Wis.-----	30	15	1	-	Long Beach, Calif.---	85	50	4	8
Milwaukee, Wis.-----	139	77	7	13	Los Angeles, Calif.---	569	359	24	28
Peoria, Ill.-----	44	26	1	3	Oakland, Calif.-----	179	105	15	7
Rockford, Ill.-----	38	25	3	3	Pasadena, Calif.-----	44	33	1	1
South Bend, Ind.-----	40	33	2	2	Portland, Oreg.-----	130	77	2	7
Toledo, Ohio-----	92	61	2	3	Sacramento, Calif.---	75	43	3	9
Youngstown, Ohio-----	59	33	2	2	San Diego, Calif.-----	108	58	4	4
					San Francisco, Calif.--	209	107	6	5
WEST NORTH CENTRAL:	907	577	28	43	San Jose, Calif.-----	41	26	4	2
Des Moines, Iowa-----	62	38	2	2	Seattle, Wash.-----	147	92	4	8
Duluth, Minn.-----	16	13	-	1	Spokane, Wash.-----	48	32	1	2
Kansas City, Kans.---	49	26	4	5	Tacoma, Wash.-----	37	21	1	1
Kansas City, Mo.-----	136	81	2	9					
Lincoln, Nebr.-----	46	33	2	1	Total	13,546	7,781	638	715
Minneapolis, Minn.---	96	65	2	7	Cumulative Totals				
Omaha, Nebr.-----	77	53	4	1	including reported corrections for previous weeks				
St. Louis, Mo.-----	262	168	4	8	All Causes, All Ages -----			27,283	
St. Paul, Minn.-----	83	58	1	3	All Causes, Age 65 and over-----			15,698	
Wichita, Kans.-----	80	42	6	7	Pneumonia and Influenza, All Ages-----			1,326	
					All Causes, Under 1 Year of Age-----			1,396	

*Estimate - based on average percent of divisional total.

HEPATITIS SURVEILLANCE SUMMARY

The total cases and the reported incidence of viral hepatitis in the United States in the first two quarters of the epidemiological year 1965-66* are the lowest reported since 1959-60. The continuing decline in incidence since the 1960-61 peak year is evident in Figure 2, page 11.

There were 7,361 cases of viral hepatitis reported in the U.S. during the summer quarter and 8,150 cases reported during the fall quarter of the epidemiological year 1965-66. This represents rates of 3.8 and 4.2 cases per 100,000 population for these two quarters, respectively (Table 2).

(Reported by the Hepatitis Surveillance Unit, CDC.)

*Hepatitis morbidity data are summarized in terms of an "epidemiological year," which runs from the 27th week of each year through the 26th week of the succeeding year.

Table 2
A. NUMBER OF REPORTED CASES OF VIRAL
HEPATITIS PER QUARTER

Epidemiological year	Summer quarter	Fall quarter	Winter quarter	Spring quarter	Total year
1958-59	3,403	4,028	7,169	4,902	19,502
1959-60	4,700	6,026	9,793	9,917	30,436
1960-61	8,940	12,403	23,096	19,598	65,267
1961-62	14,229	15,580	17,995	13,533	61,337
1962-63	10,272	11,330	13,848	9,864	45,314
1963-64	8,969	10,250	12,118	9,320	40,657
1964-65	7,581	8,597	10,311	4,876	34,365
1965-66	7,361	8,150

B. REPORTED CASES OF VIRAL HEPATITIS PER
100,000 POPULATION PER QUARTER

Epidemiological year	Summer quarter	Fall quarter	Winter quarter	Spring quarter	Total year
1958-59	2.0	2.3	4.0	2.8	11.1
1959-60	2.7	3.4	5.5	5.5	17.1
1960-61	5.0	6.9	12.8	11.1	35.8
1961-62	7.9	8.7	9.8	7.4	33.8
1962-63	5.6	6.2	7.4	5.3	24.5
1963-64	4.8	5.4	6.4	4.9	21.6
1964-65	4.0	4.5	5.3	4.2	18.0
1965-66	3.8	4.2

THE MORBIDITY AND MORTALITY WEEKLY REPORT, WITH A CIRCULATION OF 15,300, IS PUBLISHED AT THE COMMUNICABLE DISEASE CENTER, ATLANTA, GEORGIA.

CHIEF, COMMUNICABLE DISEASE CENTER
CHIEF, EPIDEMIOLOGY BRANCH
ACTING CHIEF, STATISTICS SECTION
CHIEF, SURVEILLANCE SECTION


JAMES L. GODDARD, M.D.
A.D. LANGMUIR, M.D.
IDA L. SHERMAN, M.S.
D.A. HENDERSON, M.D.
D.J.M. MCKENZIE, M.B.,
F.R.C.P.E.

IN ADDITION TO THE ESTABLISHED PROCEDURES FOR REPORTING MORBIDITY AND MORTALITY, THE COMMUNICABLE DISEASE CENTER WELCOMES ACCOUNTS OF INTERESTING OUTBREAKS OR CASE INVESTIGATIONS WHICH ARE OF CURRENT INTEREST TO HEALTH OFFICIALS AND WHICH ARE DIRECTLY RELATED TO THE CONTROL OF COMMUNICABLE DISEASES. SUCH COMMUNICATIONS SHOULD BE ADDRESSED TO:

THE EDITOR
MORBIDITY AND MORTALITY WEEKLY REPORT
COMMUNICABLE DISEASE CENTER
ATLANTA, GEORGIA 30333

NOTE: THE DATA IN THIS REPORT ARE PROVISIONAL AND ARE BASED ON WEEKLY TELEGRAMS TO THE CDC BY THE INDIVIDUAL STATE HEALTH DEPARTMENTS. THE REPORTING WEEK CONCLUDES ON SATURDAY; COMPILED DATA ON A NATIONAL BASIS ARE RELEASED ON THE SUCCEEDING FRIDAY.

U. S. DEPARTMENT OF
HEALTH, EDUCATION, AND WELFARE
Public Health Service
Communicable Disease Center
Atlanta, Georgia 30333
Official Business



POSTAGE AND FEES PAID
U. S. DEPARTMENT OF H. E. W.

ERRATUM, Vol. 14, No. 52, p. 444

Current Trends - Poliomyelitis

The isolation of virus from the case in Michigan should read "Type III poliovirus and Coxsackie virus Type A4 were also isolated from one specimen."